

Patient Name:

DOB:

Orencia® (abatacept) Infusion Orders

Psoriatic Arthritis	Rheumatoid Arthritis	Other:
(ICD-10) (I	CD-10)	(ICD-10)
Nursing Orders:		
 Hold infusion and notify p 		
• • •	s of illness or active infectior	1
o Planned/recent su	•	
o Recent live vaccin		
Record vital signs before		
Monitor and record weig		
		ollow Hypersensitivity Reaction
Management Protocol as	clinically indicated.	
Premedications: Tylenol 500 mg PO PRN	U oratadina 10 mg PO P	RN Solu-Medrol 125 mg IVP
Benadryl 25 mg PO PRN	IOther:	
using a sterile, non-pyrogenic, lo	•	e size 0.2 to 1.2 microns).
using a sterile, non-pyrogenic, lo Current w	w protein-binding filter (pore	
using a sterile, non-pyrogenic, lo Current w (Recommended do	w protein-binding filter (pore	e size 0.2 to 1.2 microns).
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one):	w protein-binding filter (pore	e size 0.2 to 1.2 microns). 5 kg: 750 mg; >100 kg: 1000 mg)
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one):	w protein-binding filter (pore eight: kg ose range: < 60 kg: 500 mg; 60-100	e size 0.2 to 1.2 microns). 5 9 kg: 750 mg; >100 kg: 1000 mg)
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one): On Week 0, Week	w protein-binding filter (pore eight: kg ose range: < 60 kg: 500 mg; 60-100 2, Week 4, then every 4 wee	e size 0.2 to 1.2 microns). 5 kg: 750 mg; >100 kg: 1000 mg)
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one):	w protein-binding filter (pore eight: kg ose range: < 60 kg: 500 mg; 60-100 2, Week 4, then every 4 wee	e size 0.2 to 1.2 microns). 5 9 kg: 750 mg; >100 kg: 1000 mg)
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one): On Week 0, Week Every 4 weeks	w protein-binding filter (pore eight: kg ose range: < 60 kg: 500 mg; 60-100 2, Week 4, then every 4 wee	e size 0.2 to 1.2 microns). 5 9 kg: 750 mg; >100 kg: 1000 mg)
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one): On Week 0, Week Every 4 weeks	w protein-binding filter (pore eight: kg ose range: < 60 kg: 500 mg; 60-100 2, Week 4, then every 4 wee	e size 0.2 to 1.2 microns). 5 9 kg: 750 mg; >100 kg: 1000 mg)
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one): On Week 0, Week Every 4 weeks	w protein-binding filter (pore eight: kg ose range: < 60 kg: 500 mg; 60-100 2, Week 4, then every 4 wee	e size 0.2 to 1.2 microns). 5 9 kg: 750 mg; >100 kg: 1000 mg)
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one): On Week 0, Week	w protein-binding filter (pore eight: kg ose range: < 60 kg: 500 mg; 60-100 2, Week 4, then every 4 wee	e size 0.2 to 1.2 microns). 5 9 kg: 750 mg; >100 kg: 1000 mg)
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one): On Week 0, Week Every 4 weeks	w protein-binding filter (pore eight: kg ose range: < 60 kg: 500 mg; 60-100 2, Week 4, then every 4 wee	e size 0.2 to 1.2 microns). 5 9 kg: 750 mg; >100 kg: 1000 mg)
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one): On Week 0, Week Every 4 weeks Every 4 weeks Everyv Additional Orders:	w protein-binding filter (pore eight: kg ose range: < 60 kg: 500 mg; 60-100 2, Week 4, then every 4 wee weeks	e size 0.2 to 1.2 microns). kg: 750 mg; >100 kg: 1000 mg) ks
using a sterile, non-pyrogenic, lo Current w (Recommended do Frequency (chose one): On Week 0, Week Every 4 weeks Every 4 weeks Everyv Additional Orders: Provider (please print):	w protein-binding filter (pore eight: kg ose range: < 60 kg: 500 mg; 60-100 2, Week 4, then every 4 wee weeks	e size 0.2 to 1.2 microns). kg: 750 mg; >100 kg: 1000 mg) ks

per manufacturer's instructions as necessitated by product availability.