



**LOUIS H.
MEDVED, MD**
INFUSION THERAPY

Patient Name: _____
DOB: _____
Date of last treatment: _____

VYEPTI® (eptinezumab-jjmr) Infusion Orders

Diagnosis (please provide ICD-10 code in space provided):

_____ Migraine
(ICD-10)

_____ Other: _____
(ICD-10)

- Hold infusion and notify provider for:
 - Abnormal vital signs
 - History of hypersensitivity to VYEPTI
 - Chance of pregnancy
- If infusion-related reaction occurs, stop infusion, notify provider and follow Hypersensitivity Reaction Management Protocol as clinically indicated.

- Administer VYEPTI® eptinezumab-jjmr 100 mg in 100 mL 0.9% sodium chloride over a period of 30 minutes. Flush with 20 ml 0.9% sodium chloride following infusion.
- Administer VYEPTI® eptinezumab-jjmr 300mg in 100 mL 0.9% sodium chloride over a period of 30 minutes. Flush with 20 mL 0.9% sodium chloride following infusion.

Use a 0.2 to 0.22micron in-line or add-on sterile filter.

Frequency (chose one):

Every 3 months

Other _____

Additional Orders:

Ordering Provider: (Louis Medved)

(Other): _____

Provider signature: _____ Date: _____